

How RAZADYNE™ ER (galantamine HBr) Can Help

A father getting lost on his way to the golf course he's played for years. A mother forgetting the recipe she's known by heart since her children were small. When you know something is wrong, once-daily RAZADYNE ER can help someone with Alzheimer's disease (AD) stay engaged in daily activities.

- In six-month clinical studies, RAZADYNE ER helped many patients maintain their cognitive (memory and thinking) and functional abilities (ability to accomplish everyday tasks)³.
- Helps patients help themselves.
- RAZADYNE ER is taken just once a day, so it's convenient for you and your loved one.
- Please see Important Product Information for RAZADYNE ER.
- Talk to your healthcare professional or visit www.RazadyneER.com for more information about how RAZADYNE ER can help.

Please see accompanying full Prescribing Information for RAZADYNE ER.

References:

1. Alzheimer's Association. Basics of Alzheimer's disease: what it is and what can you do. Chicago, IL; 2005: 24.
2. Danysz W, Parsons CG, Mobius HJ, Stoffler A, Quack G. Neuroprotective and symptomatological action of memantine relevant for Alzheimer's disease: a unified glutamatergic hypothesis on the mechanism of action. *Neurotoxicity Res.* 2000;2:85-97.
3. Brodaty H, Corey-Bloom J, Potocnik FCV, Truyen L, Gold M, Damaraju CRV. Galantamine prolonged-release formulation in the treatment of mild to moderate Alzheimer's disease. *Dement Geriatr Cogn Disord.* 2005;20:120-132.



RAZADYNE ER is used to treat the symptoms of Alzheimer's disease.

Important Safety Information

In clinical trials, the most frequent adverse events with RAZADYNE ER were similar to the twice-daily formulation of galantamine HBr. With the twice-daily formulation of galantamine HBr, the most common side effects were nausea, vomiting, diarrhea, loss of appetite and weight loss. Typically, these side effects are mild and temporary.

RAZADYNE ER may not be for everyone. Some people experience a slowed heart rate, which may lead to fainting. Serious stomach problems can occur in people taking medications such as nonsteroidal anti-inflammatory drugs (NSAIDs) or those at risk for stomach ulcers.

Compliments of



Together Rx Access is a trademark of Together Rx Access, LLC.